

Central Venous Catheter-Related Blood Stream Infection Rate

Patient Safety Indicators #23 Technical Specifications

Area-Level Indicator

AHRQ Quality Indicators, Version 4.3, August 2011

Version 4.3a is a maintenance release of Version 4.3. The differences between the two versions are:

- Version 4.3a includes Version 29 of the Limited License edition of the 3M™ APR-DRG Grouper. This grouper corrects an issue found in Version 28 of the Limited License edition of the 3M™ APR-DRG Grouper. The grouper is only used with the Inpatient Quality Indicator (IQI) mortality measures. Version 4.3 includes Version 28 of the Limited License edition of the 3M™ APR-DRG Grouper, which was incorrectly assigning a Risk of Mortality (ROM) subclass for cases dated on or after 10/1/10.
- Version 4.3a allows users to calculate area-level indicators for years 2010 and 2011, correcting an issue previously identified in Version 4.3.

All other aspects of the software, including measure specifications, remain the same. Thus this document (related to Version 4.3) remains unchanged.

Numerator

Discharges, 18 years and older **or** MDC 14 (pregnancy, childbirth, and puerperium), with selected infections defined by specific ICD-9-CM codes in any diagnosis field among all medical and surgical discharges defined by specific DRGs or MS-DRGs.

For discharges prior to October 1, 2007:

ICD-9-CM Hospital-associated infection diagnosis codes:

99662	DUE TO OTHER VASCULAR DEVICE, IMPLANT, AND GRAFT	9993	OTHER INFECTION
-------	---	------	-----------------

For discharges on or after October 1, 2007:

ICD-9-CM Central line-associated blood stream infection diagnosis code:

99931	INFECTION DUE TO CENTRAL VENOUS CATHETER
-------	---

See *Patient Safety Indicators Appendices*:

- Appendix B – Medical Discharge DRGs
- Appendix C – Medical Discharge MS-DRGs
- Appendix D – Surgical Discharge DRGs
- Appendix E – Surgical Discharge MS-DRGs

Exclude cases:

- with any diagnosis or procedure code for immunocompromised state

- with any diagnosis of cancer
- with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing) or principal diagnosis (DX1=missing), county (PSTCO=missing)

See *Patient Safety Indicators Appendices*:

- Appendix H – Cancer Diagnosis Codes
- Appendix I – Immunocompromised State Diagnosis and Procedure Codes

Denominator

Discharges in the numerator are assigned to the denominator based on the Metro Area¹ or county of the patient residence, not the Metro Area or county of the hospital where the discharge occurred.

¹ The term "metropolitan area" (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs) and primary metropolitan statistical areas (PMSAs). In addition, "area" could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999 OMB Metropolitan Statistical Area or 3) 2003 OMB Metropolitan Statistical Area. Micropolitan Statistical Areas are not used in the QI software.