

Prevention Quality Indicator 12 (PQI 12) Urinary Tract Infection Admission Rate July 2019 Area-Level Indicator Type of Score: Rate

Prepared by:

Agency for Healthcare Research and Quality U.S. Department of Health and Human Services www.qualityindicators.ahrq.gov

DESCRIPTION

Admissions with a principal diagnosis of urinary tract infection per 100,000 population, ages 18 years and older. Excludes kidney or urinary tract disorder admissions, other indications of immunocompromised state admissions, obstetric admissions, and transfers from other institutions.

[NOTE: The software provides the rate per population. However, common practice reports the measure as per 100,000 population. The user must multiply the rate obtained from the software by 100,000 to report admissions per 100,000 population.]

NUMERATOR

Discharges, for patients ages 18 years and older, with a principal ICD-10-CM diagnosis code for urinary tract infection (*ACSUTID**).

[NOTE: Obstetric discharges are not included in the PQI rate for PQI 12, though the AHRQ QI[™] software does not explicitly exclude obstetric cases. By definition, discharges with a principal diagnosis of urinary tract infection exclude obstetric discharges, because the principle diagnosis for an obstetric discharge would identify it as obstetric, and no such diagnoses are included in the set of qualifying diagnoses.]

NUMERATOR EXCLUSIONS

Exclude cases:

• with admission source for transferred from a different hospital or other health care facility (*Appendix A*) (UB04 Admission source - 2, 3)

• with a point of origin code for transfer from a hospital, skilled nursing facility (SNF) or intermediate care facility (ICF), or other healthcare facility (*Appendix A*) (UB04 Point of Origin - 4, 5, 6)

• with any-listed ICD-10-CM diagnosis codes for kidney/urinary tract disorder (*KIDNEY**)

• with any-listed ICD-10-CM diagnosis codes (Appendix C: IMMUNID) or any-listed ICD-10-PCS

procedure codes for immunocompromised state ((Appendix C: IMMUNIP)

• with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year

(YEAR=missing), principal diagnosis (DX1=missing), or county (PSTCO=missing)

<u>Appendix A – Admission Codes for Transfers</u> Appendix C - Immunocompromised State Diagnosis and Procedure Codes

DENOMINATOR

Population ages 18 years and older in metropolitan area^{**} or county. Discharges in the numerator are assigned to the denominator based on the metropolitan area or county of the patient residence, not the metropolitan area or county of the hospital where the discharge occurred.

** The term "metropolitan area" (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs), and primary metropolitan statistical areas (PMSAs). In addition, "area" could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999 OMB Metropolitan Statistical Area, or 4) 2003 OMB Metropolitan Statistical Area. Micropolitan Statistical Areas are not used in the QI software.

* See below for code list

AHRQ QI[™] ICD-10-CM/PCS Specification v2018 PQI 12 Urinary Tract Infection Admission Rate www.qualityindicators.ahrq.gov

Urinary tract infection diagnosis codes: (ACSUTID)

N10 N12	Acute tubulo-interstitial nephritis Tubulo-interstitial nephritis, not specified as acute or chronic	N2886 N3000	Ureteritis cystica Acute cystitis without hematuria
N151	Renal and perinephric abscess	N3001	Acute cystitis with hematuria
N159	Renal tubulo-interstitial disease, unspecified	N3090	Cystitis, unspecified without hematuria
N16	Renal tubulo-interstitial disorders in diseases classified elsewhere	N3091	Cystitis, unspecified with hematuria
N2884	Pyelitis cystica	N390	Urinary tract infection, site not specified

N2885 Pyeloureteritis cystica

July 2	2019
--------	------

Kidney or urinary tract disorder diagnosis codes: (KIDNEY)

N110	Nonobstructive reflux-associated chronic pyelonephritis	Q6231	Congenital ureterocele, orthotopic
N111	Chronic obstructive pyelonephritis	Q6232	Cecoureterocele
N118	Other chronic tubulo-interstitial nephritis	Q6239	Other obstructive defects of renal pelvis and
		C	ureter
N119	Chronic tubulo-interstitial nephritis, unspecifie	d Q624	Agenesis of ureter
N130	Hydronephrosis with ureteropelvic junction	Q625	Duplication of ureter
	obstruction		*
N136	Pyonephrosis	Q6260	Malposition of ureter, unspecified
N1370	Vesicoureteral-reflux, unspecified	Q6261	Deviation of ureter
N1371	Vesicoureteral-reflux without reflux nephropathy	Q6262	Displacement of ureter
N13721	Vesicoureteral-reflux with reflux nephropathy	Q6263	Anomalous implantation of ureter
110,21	without hydroureter, unilateral	20200	
N13722	Vesicoureteral-reflux with reflux nephropathy	Q6269	Other malposition of ureter
	without hydroureter, bilateral		I I
N13729	Vesicoureteral-reflux with reflux nephropathy	Q627	Congenital vesico-uretero-renal reflux
	without hydroureter, unspecified		-
N13731	Vesicoureteral-reflux with reflux nephropathy	Q628	Other congenital malformations of ureter
	with hydroureter, unilateral		
N13732	Vesicoureteral-reflux with reflux nephropathy	Q630	Accessory kidney
	with hydroureter, bilateral		
N13739	Vesicoureteral-reflux with reflux nephropathy	Q631	Lobulated, fused and horseshoe kidney
N120	with hydroureter, unspecified	0(22	Esteria bidu en
N139	Obstructive and reflux uropathy, unspecified	Q632	Ectopic kidney
Q600	Renal agenesis, unilateral	Q633	Hyperplastic and giant kidney
Q601	Renal agenesis, bilateral	Q638	Other specified congenital malformations of
			kidney
Q602	Renal agenesis, unspecified	Q639	Congenital malformation of kidney,
			unspecified
Q603	Renal hypoplasia, unilateral	Q6410	Exstrophy of urinary bladder, unspecified
Q604	Renal hypoplasia, bilateral	Q6411	Supravesical fissure of urinary bladder
Q605	Renal hypoplasia, unspecified	Q6412	Cloacal extrophy of urinary bladder
Q606	Potter's syndrome	Q6419	Other exstrophy of urinary bladder
Q6100	Congenital renal cyst, unspecified	Q642	Congenital posterior urethral valves
Q6101	Congenital single renal cyst	Q6431	Congenital bladder neck obstruction
Q6102	Congenital multiple renal cysts	Q6432	Congenital stricture of urethra
Q6111	Cystic dilatation of collecting ducts	Q6433	Congenital stricture of urinary meatus
Q6119	Other polycystic kidney, infantile type	Q6439	Other atresia and stenosis of urethra and
			bladder neck
Q612	Polycystic kidney, adult type	Q645	Congenital absence of bladder and urethra
Q613	Polycystic kidney, unspecified	Q646	Congenital diverticulum of bladder
Q613 Q614	Renal dysplasia	Q6470	Unspecified congenital malformation of
X014	Konai ayspiasia	v ⁰⁺⁷⁰	bladder and urethra
Q615	Medullary cystic kidney	Q6471	Congenital prolapse of urethra
×010		-	
		luly 2019	

AHRQ QI[™] ICD-10-CM/PCS Specification v2019 PQI 12 Urinary Tract Infection Admission Rate www.qualityindicators.ahrq.gov

Q618	Other cystic kidney diseases	Q6472	Congenital prolapse of urinary meatus
Q619	Cystic kidney disease, unspecified	Q6473	Congenital urethrorectal fistula
Q620	Congenital hydronephrosis	Q6474	Double urethra
Q6210	Congenital occlusion of ureter, unspecified	Q6475	Double urinary meatus
Q6211	Congenital occlusion of ureteropelvic junction	Q6479	Other congenital malformations of bladder and urethra
Q6212	Congenital occlusion of ureterovesical orifice	Q648	Other specified congenital malformations of urinary system
Q622	Congenital megaureter	Q649	Congenital malformation of urinary system, unspecified